

# 1 Coral reef fish populations can persist without immigration

## 2 Running Title: Reef fish population persistence

3 Océane C. Salles<sup>1,\*</sup>, Jeffrey A. Maynard<sup>1,2</sup>, Marc Joannides<sup>3</sup>, Corentin M. Barbu<sup>4</sup>, Pablo Saenz-  
4 Agudelo<sup>5</sup>, Glenn R. Almany<sup>1</sup>, Michael L. Berumen<sup>6</sup>, Simon R. Thorrold<sup>7</sup>, Geoffrey P. Jones<sup>8</sup>, and  
5 Serge Planes<sup>1</sup>

6 *1- Laboratoire d'Excellence "CORAIL", USR 3278 CNRS-EPHE-UPVD CRIOBE, Perpignan,*  
7 *France.*

8 *2- SymbioSeas and Marine Applied Research Center, Wilmington, NC 28411, United States of*  
9 *America.*

10 *3- Université de Montpellier 2, UMR 5149 I3M, Pl Eugene Bataillon, F-34095 Montpellier,*  
11 *France.*

12 *4- Department of Biostatistics and Epidemiology, Center for Clinical Epidemiology and*  
13 *Biostatistics, University of Pennsylvania, 800 Blockley Hall, Philadelphia, PA 19104, United States*  
14 *of America.*

15 *5- Instituto de Ciencias Ambientales y Evolutivas, Universidad Austral de Chile, Chile.*

16 *6- Red Sea Research Center, Division of Biological and Environmental Sciences and Engineering,*  
17 *King Abdullah University of Science and Technology, Thuwal, 23955, Saudi Arabia.*

18 *7- Biology Department, Woods Hole Oceanographic Institution, Woods Hole, MA 02543, United*  
19 *States of America.*

20 *8- ARC Centre of Excellence for Coral Reef Studies and School of Marine and Tropical Biology,*  
21 *James Cook University, Townsville, Queensland 4811, Australia.*

22 \* Author for correspondence: Océane C. Salles (oceane.salles@gmail.com)

## 23 Abstract

24 Determining the conditions under which populations may persist requires accurate estimates of  
25 demographic parameters, including immigration, local reproductive success, and mortality rates. In  
26 marine populations, empirical estimates of these parameters are rare, due at least in part to the  
27 pelagic dispersal stage common to most marine organisms. Here, we evaluate population  
28 persistence and turnover for a population of orange clownfish, *Amphiprion percula*, at Kimbe Island  
29 in Papua New Guinea. All fish in the population were sampled and genotyped on five occasions at  
30 2-year intervals spanning eight years. The genetic data enabled estimates of reproductive success  
31 retained in the same population (reproductive success to self-recruitment), reproductive success  
32 exported to other subpopulations (reproductive success to local connectivity), and immigration and  
33 mortality rates of sub-adults and adults. Approximately 50% of the recruits were assigned to parents  
34 from the Kimbe Island population and this was stable through the sampling period. Stability in the  
35 proportion of local and immigrant settlers is likely due to: low annual mortality rates and stable egg  
36 production rates, and the short larval stages and sensory capacities of reef fish larvae. Biannual  
37 mortality rates ranged from 0.09 to 0.55 and varied significantly spatially. We used these data to  
38 parameterize a model that estimated the probability of the Kimbe Island population persisting in the  
39 absence of immigration. The Kimbe Island population was found to persist without significant  
40 immigration. Model results suggest the island population persists because the largest of the  
41 subpopulations are maintained due to having low mortality and high self-recruitment rates. Our  
42 results enable managers to appropriately target and scale actions to maximize persistence likelihood  
43 as disturbance frequencies increase.

44 **Keywords:** *Amphiprion percula* | long-term monitoring | parentage analysis | self-recruitment |  
45 population demography | persistence

## 1. Introduction

Understanding and identifying biotic and abiotic conditions under which wild populations can persist over space and through time is a formidable theoretical and practical challenge in population ecology [1,2]. Identifying the spatial scale at which populations are influenced by these conditions allows us to define conservation management units and implement appropriately targeted and scaled actions. Wild populations of many species are fragmented and function as metapopulations, comprised of local populations linked by migration [3]. Within metapopulations, local population size is driven by four fundamental demographic rates: recruitment and immigration, which increase the population size, as well as mortality and emigration, which decrease the population size [4]. Local populations within metapopulations can only persist when recruitment and immigration equals or exceeds mortality and emigration [5,6]. However, these demographic rates may equalize as populations approach carrying capacity, a theoretical upper limit on the population size [7,8].

Population models provide a useful way of examining the influence of demographic parameters on population persistence [2,9,10]. These models can be used to generate projections of how the size of a population may change under different environmental conditions, movement patterns of individuals within populations, and mortality rates. Demographic models may also include some level of density dependence, although such processes are hard to estimate in wild populations [11]. However, the reliability of demographic models depends on the accuracy of parameters that are often estimated indirectly due to logistical sampling constraints [12–14]. Evaluating demographic parameters is challenging because it is often difficult or impossible to follow a large number of individuals in situ over a time period sufficiently long enough to accurately estimate rates of population turnover [15]. This is particularly the case for organisms that are not sessile, with complex life cycles that include dispersal, ontogenetic movements, hibernation, or cryptic phases.

70 Building demographic models is especially challenging for marine species as they often have  
 71 complex life cycles that may include a dispersive larval phase and/or a mobile adult phase.  
 72 Consequently, it is often difficult to distinguish between losses due to emigration and losses due to  
 73 mortality [16]. Many studies have been forced to assume that emigration equals immigration, but it  
 74 is then impossible to quantify the relative contributions of these two processes to the dynamics of  
 75 local populations within metapopulations [17,18]. This assumption can bias models towards  
 76 overestimating the contribution of local populations to the persistence of the metapopulation.  
 77 Identifying the natal origins of larvae recruiting to marine populations has also represented a  
 78 significant challenge to the development of accurate demographic models. Larval recruitment to  
 79 subpopulations comes from two potential sources. Individuals may return to their natal  
 80 subpopulation (typically termed self-recruitment), or may disperse to other subpopulations within  
 81 the metapopulation (typically termed connectivity). Both self-recruitment and connectivity may be  
 82 critical for the persistence of subpopulations, but the means to determine their relative importance  
 83 has not been available until recently.

84 The development of several methods to determine the natal origins of juveniles has provided new  
 85 insights into the spatial scale of larval dispersal (e.g. physical oceanography [19,20], artificial  
 86 tagging [10], trace-elemental fingerprinting of fish otoliths [21] and invertebrate larval shells [22],  
 87 and molecular genetics [23,24]). For instance, parentage analysis based on DNA genotyping  
 88 provides a means to distinguish natal origins as it allows for detection of parent–child relationships  
 89 [24]. We have used this approach to estimate larval dispersal of orange clownfish in Kimbe Bay  
 90 from samples collected in 2005 and 2007 [24,25]. When applied regularly to all individuals in a  
 91 population, DNA fingerprinting using microsatellite loci also provides a powerful tool for  
 92 measuring mortality and population turnover. Orange clownfish (*Amphiprion percula*) form  
 93 obligate associations with sea anemones, where they remain for the rest of their lives. Mortality and  
 94 population turnover rates can be quantified by tracking individuals from each subpopulation  
 95 through time. Previous results from such studies showed a high self-recruitment rate (approx. 50%

of recruitment) for the *A. percula* of Kimbe Island in 2005 and 2007 [24,25] and a very low adult mortality rate (approx. 12.9% per year, [26]) for the same species in Madang, Papua New Guinea. Based on these findings, we hypothesized that levels of self-recruitment may be high enough to ensure the self-persistence of the *A. percula* population in Kimbe Island without immigration. We test this hypothesis using parentage analysis and DNA identification of all of the individuals in the orange clownfish population in Kimbe Island, Papua New Guinea over an 8-year sampling period. We used the resulting demographic data to parameterize a demographic model of Kimbe Island subpopulations to address a specific question: can the orange clownfish population of Kimbe Island and any of its subpopulations persist in the absence of larval connectivity from other locations in the region?

## 2. Material and methods

### (a) *The study system*

*Amphiprion percula* typically lives in groups and occurs in obligate association with sea anemones [27,28]. Groups are usually composed of a dominant breeding pair and between 0 and 6 juveniles[29]. Within each group, there is a size-based dominance hierarchy: the female is largest, the male is second largest and the non-breeders rank progressively lower in the hierarchy as they decrease in size. If the single female adult (i.e. rank 1) of a group dies, then the male (i.e. rank 2) changes sex and the largest juvenile (i.e. rank 3) from the anemone becomes male [30,31]. Reproduction occurs year round, and breeders hatch hundreds of eggs each lunar month but the importance of the lunar cycle on larval settlement is poorly understood [32,33]. Then, eggs hatch into pelagic larvae that spend approximately two weeks in the pelagic environment before settling on an anemone [10,24,34], which may or may not be on the natal reef. Because adult clownfish rarely move among anemones, persistence at the scale of a single anemone (and hence the larger scale) is determined by whether the rate of settling larvae can offset the mortality of juveniles and adults.

121 **(b) Study site and field collections**

122 The study was conducted every two years from 2005 to 2013 during April in Kimbe Island  
 123 (5°12'22.56" S, 150°22'35.58" E), a small island located in Kimbe Bay, Papua New Guinea  
 124 (electronic supplementary material, figure S1). We subdivided the island into seven geographical  
 125 areas (A–G). In this study, these geographical areas are described as subpopulations. We located  
 126 and tagged a total of 210 anemones that were present during the five periods of sampling, all of  
 127 which were occupied by *A. percula* (electronic supplementary material, figure S1). All fish at each  
 128 anemone were captured by divers using hand nets and processed in situ. Individuals were measured  
 129 using calipers (total length TL, mm), fin-clipped for genetic analysis and then released back on the  
 130 same anemone. The biggest fish in each anemone was identified as the female, the second largest  
 131 individual was assumed to be the male and all other individuals were classified as juveniles. In 2005  
 132 and 2007, juveniles (non-breeders) with TL > 35 mm were not fin-clipped but were recorded for  
 133 2007. The small pieces of fin tissue were preserved in 95% ethanol in 0.2 ml vials. The smallest  
 134 non-breeders (TL < 35 mm, hereafter referred to as ‘new-recruits’) were removed from their  
 135 anemones and the whole fish were preserved in vials with 95% ethanol.

136 Resident clownfish may prevent recruitment of new individuals at high densities [32]. Therefore,  
 137 removal of the small individuals homogenized conditions for larval settlement among all anemones.  
 138 All details related to capture numbers for each stage, subpopulation and year of sampling are  
 139 provided within the electronic supplementary material, table S1. DNA from a total of 3 890  
 140 collected samples (including 1 067 new-recruits) were extracted and screened at 22 polymorphic  
 141 microsatellite loci (electronic supplementary material, table S2) that satisfied Hardy–Weinberg  
 142 equilibrium and linkage disequilibrium assumptions.

143 **(c) Demographic parameters**

144 *Mortality measurement (biannual).*

145 The package GenAlex v6.5 [35] was used to compare each of the individual genotypes from fish  
 146 sampled in 2009, 2011, and 2013 (2005 and 2007 are excluded because juveniles with TL > 35 mm  
 147 were not fin-clipped in those years) to identify all unique multilocus genotypes [36]. The combined  
 148 probability of identity (i.e. the probability of two samples having the same multilocus genotype by  
 149 chance) for this set of markers in this population was small ( $1.7 \times 10^{-24}$ ). Therefore, if fin clips from  
 150 different sampling years had the same genotype, they were assumed to be from the same individual.  
 151 To account for possible genotyping errors, mismatches of up to two loci between genotypes were  
 152 allowed in pair-wise genotype comparisons. In this way, we were able to measure mortality for each  
 153 stage (i.e. average mortality of females, males, and juveniles) for each of the seven subpopulations  
 154 of Kimbe Island for two time steps (i.e. 2009–2011 and 2011–2013). We used  $\chi^2$ -test to compare  
 155 the mortality rates observed among the seven subpopulations and between the various life stages  
 156 (i.e. juvenile, male, and female stages). We then used the Fisher-test to evaluate whether the  
 157 mortality rate calculated for 2009–2011 was significantly different than the mortality rate calculated  
 158 for 2011–2013.

159 *Reproductive success to self-recruitment and to local connectivity, and immigration measurements*  
 160 *(biannual).*

161 For every sampling year (2005, 2007, 2009, 2011, and 2013) and for each subpopulation of Kimbe  
 162 Island (A–G), we used assignment/exclusion tests to estimate an index of reproductive success to  
 163 self-recruitment, reproductive success to local connectivity, and immigration. Reproductive success  
 164 to self-recruitment is calculated as the number of juveniles with parents from the same  
 165 subpopulation divided by the number of females in this population. Reproductive success to local  
 166 connectivity is calculated as the number of juveniles in a subpopulation that were offspring of  
 167 parents from another Kimbe Island subpopulation, divided by the number of females in the focal  
 168 population. Immigration is defined as the number of juveniles that were not assigned to any parent  
 169 from Kimbe Island. To assign one individual to a breeding pair, we used individual genotypes and  
 170 the platform FaMoz [37]. We then compared the percentages of self- and local-recruits and

immigrants within Kimbe Island among all sampling years using a Kruskal–Wallis test. In this analysis, local-recruitment is the percentage of self-recruitment, defined by Burgess et al. [15] as the fraction of recruitment to a site comprised of individuals born in that site and the percentage of local connectivity. We used the Pettitt homogeneity test to evaluate differences in the presence of connections among subpopulations (i.e. exchange of recruits) and the number of recruits being exchanged among (local recruits) and within (self-recruits) subpopulations among years.

#### *Carrying capacity.*

The number of female and male clownfish per anemone cannot exceed one since anemones only host one breeding pair. Therefore, the carrying capacity for adult males and females in a given subpopulation cannot exceed the number of anemones present in that subpopulation. The number of juveniles per anemone may vary between the different subpopulations. We calculated the mean number of clownfish observed per anemone in the seven subpopulations at four sampling periods (2007, 2009, 2011, and 2013, excluding 2005 because non-breeders more than 35 mm were not counted and sampled). For each subpopulation, we obtained a set of four values of mean number of clownfish per anemone and we used the highest values of mean number of clownfish per anemone as the juvenile carrying capacity per anemone. The juvenile carrying capacity per subpopulation is equal to the maximum number of clownfish observed per anemone multiplied by the total number of anemones in that subpopulation. Mortality rates, and rates for reproductive success to self-recruitment, reproductive success to local connectivity, and immigration are exact measures from the field observations (rather estimation) so are not presented with error bars. All statistical analyses were performed using R v3.0.2 [38] and significance was interpreted using a p-value of >0.05.

#### *(d) Demographic model*

The system is based on a loss-input principle using recurrence equations [39]. The number of larvae, juveniles, males, and females in subpopulation at period  $t + h$  (with the time step resolution of the dynamic) is by  $L_{t+h}^i, J_{t+h}^i, M_{t+h}^i$  and  $F_{t+h}^i$ . The dynamics of each subpopulation



196 are described using these four recurrence equations:

$$L_{t+h}^i = h \left[ b^{ii} w_t^i + \sum_{j=1}^{sP} b^{ji} w_t^j + \theta^i \right], \quad (2.1)$$

$$J_{t+h}^i - J_t^i = h [(-d_j^i - t_M) J_t^i + s_L L_t^i], \quad (2.2)$$

$$M_{t+h}^i - M_t^i = h [(-d_M^i - t_F) M_t^i + t_M J_t^i], \quad (2.3)$$

$$F_{t+h}^i - F_t^i = h [-d_F^i F_t^i + t_F M_t^i], \quad (2.4)$$

197 where  $w_t^i$  and  $w_t^j$  are the number of breeding pair in the subpopulation  $i$  and  $j$  at time  $t$  and (eq.  
 198 2.1),  $d_j^i$ ,  $d_M^i$  and  $d_F^i$  are the mortality rates per capita and by unit of time of juveniles, males and  
 199 females in subpopulation  $i$  (eq. 2.2 to 2.4). The parameter  $b^{ii}$  represents the number per unit of time  
 200 of self-recruits and breeding pair ( $w_t^i$ ) in subpopulation  $i$  (i.e., reproductive success to self-  
 201 recruitment), while  $b^{ji}$  is the number per unit of time of juveniles produced per breeding pair in  
 202 subpopulation  $j$  ( $w_t^j$ ) at time  $t$  that disperse and recruit into subpopulation  $i$  (i.e. reproductive  
 203 success to local connectivity).  $\sum_{j=1}^{sP} b^{ji} w_t^j$  is the sum of the connectivity from the six other  
 204 subpopulations ( $sP=6$ )  $j$  to subpopulation  $i$ . The parameter  $\theta^i$  denotes the constant number per unit  
 205 of time of migrants recruiting to subpopulation  $i$  who were spawned outside of Kimbe Island. In  
 206 equation 2.2,  $s_L$  corresponds to settlement rate per unit of time of larvae arriving at subpopulation  $i$ .  
 207 The number of larvae that can enter the juvenile population on reef  $i$  is limited by the amount of  
 208 free space,  $(n^i K^i - J_t^i)$ :

$$s_L = s_l \frac{n^i K^i - J_t^i}{n^i K^i} \quad (2.5)$$

209 where  $n^i$  is the maximal number of juveniles per anemone in subpopulation  $i$  and  $K^i$  is the number  
 210 of anemones in the subpopulation  $i$ .

211 In equations 2.2 and 2.3,  $t_M$  measures the rate of transition function from juveniles to males per unit  
 212 of time:

$$t_M = t_m \frac{K^i - M_t^i}{K^i} \quad (2.6)$$

213 where  $(K^i - M_t^i)$  denotes the space available space for new males (i.e., male mortality or males  
 214 changing into females following female mortality) taking into account that we cannot observed  
 215 more than one male per anemone.

216 In equations 2.3 and 2.4,  $t_F$  corresponds to transition function from males to females per unit of  
 217 time:

$$t_F = t_f \frac{K^i - F_t^i}{K^i} \quad (2.7)$$

218 where  $K^i - F_t^i$  denotes the female turnover due to available space for females in anemones due to  
 219 female mortality taking into account that we cannot observe more than one adult female per  
 220 anemone.

#### 221 **(e) Hypotheses, model fitting, and simulations**

222 Rates of mortality, self-recruitment, and local connectivity were all assumed to be known  
 223 parameters, deduced from the parentage analysis, which was entirely independent of the model. The  
 224 parameters  $s_i$ ,  $t_m$ , and  $t_f$  in equations (2.5)–(2.7) appear as adjustment parameters that fit the model  
 225 to the observed data (electronic supplementary material, figure S2). Their estimated values are  
 226 obtained by minimizing the least-square criterion measuring the distance between observed data  
 227 and their values predicted by the model [40], using the procedure outlined by Nelder & Mead [41].  
 228 Reproductive success rates of self- and local-recruitment, and immigration used in the demographic  
 229 model were the averages among all five sampling periods for each subpopulation (table 1).  
 230 Mortality rates used as inputs to the demography model are the averages between 2009–2011 and

2011–2013 for each life stage of each subpopulation (table 1). Demographic parameters are rescaled to equivalent monthly rates to increase resolution of the model dynamics. The model was initialized from 2007 with a monthly time step. The population dynamics were evaluated considering three different recruitment and immigration scenarios: (i) with all inputs; (ii) removing immigration; and (iii) removing local recruitment and immigration. The numerical solution for the demographic model was programmed into R v3.0.2 [38].

#### *(f) Sensitivity analysis*

Sensitivity analyses were used to determine which demographic parameters most greatly influence population dynamics and persistence among: mortality rate of females, males, and juveniles, self-recruitment and local-recruitment rates, immigration, and juvenile and adult carrying capacity. We independently varied each parameter by 30% and plotted the resulting trajectory of population size through time.

Further details on the demographic parameters and sensitivity analysis (figure S2b,f) are provided in the electronic supplementary material.

### **3. Results**

#### *(a) Population demography*

##### *Mortality measurements (biannual).*

In total, 1 706 fins were collected and genotyped between 2009 and 2013. Genetic similarity analyses revealed that 561 individuals were observed only once, 334 individuals were found in two successive sampling years (i.e. 2009 and 2011 or 2011 and 2013) and 159 individuals survived during the three years of sampling. From these data, we calculated the mortality rate between the periods 2009–2011 and 2011–2013 (electronic supplementary material, figure S3 and table S3). We excluded two cases of survival assignments (i.e. similar genotypes with three mismatches on 44 alleles from fins clipped in two consecutive periods) because fins came from females which were sampled in different subpopulations separated by more than 350 m. During the period 2009–2011,

the mortality rates differed significantly among the seven subpopulations and for the various life stages (Chi square test,  $p < 0.001$ ). During the period 2011-2013, the mortality rates among the seven subpopulations and for the various life stages were significantly different only for males and females (Chi square test,  $p_J = 0.120$ ,  $p_{M,F} < 0.001$ ). The average mortality rate varied among subpopulations between 0.18 and 0.49 for juveniles, 0.09 and 0.44 for males, and 0.19 and 0.55 for females (Table 1). The mortality rates were higher in subpopulation G than in the other subpopulations ( $d_G > d_D > d_C > d_F > d_A > d_E > d_B$ ). However, when all life stages and subpopulations were aggregated, differences in mortality rates between 2009-2011 and 2011-2013 were not significantly different (Fisher test,  $p = 0.703$ ). Further, when the development stage and subpopulations were considered separately, differences among mortality rates between the 2 periods were only significant for males and females from subpopulation A (Fisher test,  $p < 0.001$ ).

#### Recruitment measurements (biannual).

Parentage analysis was used to establish the natal origin of all *A. percula* offspring spawned by females in the seven Kimbe Island subpopulations. The analysis quantified self-recruitment (fraction of recruitment to a subpopulation comprised of new-recruits born in that subpopulation), local connectivity (fraction of recruitment to a subpopulation comprised of new-recruits born in other Kimbe Island subpopulations), and immigration from other reefs outside Kimbe Island from 2005–2013 (figure 1). From a total of 1 067 new-recruits collected over five sampling events, less than half were immigrants (489 new-recruit immigrants, 45.8%); 222 (20.9%) were self-recruits and 356 (33.3%) were local recruits). The percentages of self-recruits, local-recruits, and immigrants among sampling years did not vary significantly among years (Kruskal–Wallis test,  $p = 0.877$ ). Further, levels of self-recruitment and local connectivity among subpopulations were all consistent among the five sample years. Differences in the presence of connections among subpopulations (i.e. exchange of recruits) and the number of recruits being exchanged among (local recruits) and within (self-recruits) subpopulations were not significantly different among years (Pettitt homogeneity test,  $p = 0.702$ ). The average reproductive success to self-recruitment varied among subpopulations

282 between 0.06 and 0.45 juveniles per female (electronic supplementary material, table S4). The  
 283 reproductive success to self-recruitment was higher in subpopulation A than in the other  
 284 subpopulations ( $b_A > b_E > b_G > b_F > b_C > b_D > b_B$ ). The average number of juveniles per  
 285 female in the subpopulation A represented ~40% of the total inputs coming from Kimbe Island.  
 286 Further details on the mortality and reproductive success results are provided in the electronic  
 287 supplementary material.

#### 288 **(b) Modeling population dynamics**

289 The demographic parameters calculated from the field data and used in the model included the  
 290 reproductive success to self-recruitment, reproductive success to local connectivity, mortality and  
 291 the carrying capacity. The carrying capacity is the average number of juveniles per anemone for  
 292 juveniles and is the number of anemones (K) for adult males and females. Although the number of  
 293 juveniles per anemone was as high as 5 (figure. S4), the average number of juveniles per anemone  
 294 varied between 1.14 and 2.19 among subpopulations (table 1). The optimum parameter values for  
 295 fitting our demographic model to the observed data were  $s_l = 0.196$  (larval settlement),  $t_m = 0.310$   
 296 (transition from juvenile to male),  $t_f = 1.950$  (transition from male to female).

297 We found that all subpopulations and the Kimbe Island population were stable and persisted when  
 298 all connections were maintained (self-recruitment, local-recruitment, and immigration, figure 2  
 299 black points). The subpopulations A, B, C, D, E, and G and the integrated Kimbe Island population  
 300 were also stable and persisted when immigration was removed (figure 2, grey points). The number  
 301 of juveniles in the subpopulation F was null after 5 years (figure 2, grey points). When immigration  
 302 was removed, the presence of both a male and female on each anemone were maintained, but the  
 303 number of recruits/juveniles was halved meaning the available habitat was no longer saturated. The  
 304 global Kimbe Island population was also stable and persisted if all local connections and  
 305 immigration were removed, though the habitat became less saturated through a 25% decline in  
 306 females. In this scenario only subpopulations A, B, and E persisted (figure 2, blue points), albeit  
 307 with reductions in juvenile densities of ~50% (A and E) and ~75% (B). The juvenile stage in the

subpopulations C, D, F, and G disappeared after ~5 years. Mortality rates for these subpopulations were all higher than for A, B, and E and thus C, D, F, and G were more dependent on local connectivity and immigration. Subpopulations C, D, F, and G were smaller than A, B, and E, which is why the overall Kimbe Island population persisted when only subpopulations A, B, and E persisted. The declines in the subpopulations C, D, F and G started as reductions in the number of juvenile followed by a decrease in the number of males and then total loss due to the absence of females (electronic supplementary material, figure S5).

A sensitivity analysis was conducted to determine which parameters most influence subpopulation dynamics. The sensitivity analysis revealed that for subpopulations C, D, and F the time of extinction stayed the same irrespective of whether the other demographic parameters are modified by 30% (electronic supplementary material, figure S6a and b). For these three subpopulations, self-recruitment, local-recruitment, and immigration were equally important. Considering the entire population of females in Kimbe Island, the mortality rate of females and the carrying capacity were the most influential demographic parameters driving persistence (electronic supplementary material, figure S6a and b). The sensitivity analysis demonstrates the robustness of our main finding: the orange clownfish population from Kimbe Island can persist without immigration. In fact, we observed the persistence of the orange clownfish population even after increasing or decreasing by 30% the different demographic parameters.

## 4. Discussion

We followed an entire population of a coral reef fish through eight years to quantify demographic parameters used in assessments of population dynamics. Our results indicate that the overall Kimbe Island population of clownfish was stable with consistent patterns of self-recruitment and larval immigration, and model results suggest that the population can persist without immigration from other sources. Model results also suggest several smaller lagoonal subpopulations within Kimbe Island were also stable and likely to persist if only populated from self-recruitment. The high level

of demographic stability demonstrated at a small spatial scale is a novel and unexpected result for a marine population that includes a dispersive larval stage. The conclusion that the population can be sustained through self-recruitment is likely to be robust for two primary reasons. Firstly, our demographic model was based on genetic identification of every individual and their progeny over an 8-year period. The primary input variables to the model were therefore based directly on sampling of all individuals in the population and parent–offspring relationships, rather than use of indirect methods as is the case in other studies [12,42–44]. Further, the study was long enough to see a near-complete population turnover, with only 18% of adults persisting over the 8-year sampling period. Secondly, mortality, self-recruitment and connectivity patterns were stable over the 8-year sampling period, suggesting that the demographic importance of self-recruitment was not a transitory phenomenon. Although larval connectivity is not necessary for the persistence of relatively undisturbed populations, it is likely to be of great demographic significance via a rescue effect [45] should extrinsic disturbances extirpate a subpopulation.

The power of longitudinal genetic monitoring of each individual was that we can distinguish between death and emigration to generate an accurate estimate of mortality. Our data indicated that a small number of adult individuals moved short distances during the study period from their anemone to another anemone within the same lagoon (approx. 2%), which meant we could adjust mortality rates accordingly and calculate the true mortality rate. We calculated average biannual mortality rates of 31–36% for juveniles, and adult males and females with a global mortality rate that was constant across two 2-year periods. There was no significant difference between the mortality rate of adults (either sex) and juveniles, though most juveniles sampled were large and positioned first in the queue for access to reproduction in the colony hierarchy (juveniles named as ‘rank 3’ by Buston [26,32] represent approximately 83% of total juveniles sampled). Buston [26,32] used visual censuses to estimate mortality for the same species in another part of Papua New Guinea (Madang), and also concluded that adults (named by the authors ‘rank 1 and 2’ to refer to female and male, respectively) and juveniles (named ‘rank 3’) had similar mortality [26]. However,

359 the mortality rate we found was higher (approx. 16.5% versus approx. 12.9%) than Buston [26,32]  
360 observed following conversion of our biannual and his monthly mortality rates to annual rates.  
361 Also, differences among subpopulations were as great as approximately 25%, all of which speaks to  
362 the value of longitudinal genetic monitoring of each individual and to the importance of local-scale  
363 differences in the environment (abiotic and/or biotic) for survival.

364 Differences among subpopulations in mortality rates probably reflect a rapid turnover as a result of  
365 intense competition for reproductive status on anemones (related to the number of fish in the queue  
366 within an anemone), which are clearly a limiting resource. Only the dominant pair of clownfish  
367 breeds and non-breeding subordinates queue for breeding positions. The highest-ranking individuals  
368 are the most likely to inherit breeding positions so competition for rank may be intense, which  
369 results in high stress for some juveniles and competition for food resources within anemones  
370 [26,30]. The extent to which competition for rank plays out probably varies among subpopulations  
371 resulting in more rapid turnover where competition is most intense, which might be the case for our  
372 subpopulations C, D, and G, where mortality rates are highest.

373 For marine populations, recruitment is often a highly stochastic process with great inter-annual  
374 variation [46,47]. In our study, longitudinal genetic monitoring of each individual allowed us to  
375 distinguish between individuals that were offspring of parents from Kimbe Island subpopulations  
376 and those that were spawned by adults on other reefs (i.e. located more than 10 km from Kimbe  
377 Island). We found that local connectivity and self-recruitment combined to provide approximately  
378 50% of recruits each year during the five sampling periods. Importantly, the stability we see  
379 through time is a result of several settlement events because new-recruits sampled are between 10  
380 and 281 days old (Salles O.C., 2015) and we are therefore integrating a number of settlement events  
381 through the year. Stability in the proportion of local and immigrant settlers over time is likely due to  
382 a combination of biological and physical factors. Population numbers are fairly consistent through  
383 time with low annual mortality rates (approx. 15%), suggesting that egg production rates were also



likely to be relatively stable when averaged over the year in question. Also, the duration of the larval stage for this species is relatively short (10–12 days on average [34]) and therefore capacity for dispersal will be significantly less than for species with longer pelagic larval durations. Larval behaviour may also play a role in stabilizing self- and local-recruitment over time. Coral reef fish larvae are now known to have well-developed sensory systems and reef sounds [48–50] and olfactory cues [51–53] may be used as means to locate and orient towards home to a suitable habitat [54,55]. For instance, Dixon *et al.* [56] provided evidence that *A. percula* use olfactory cues to locate their natal reefs, providing a mechanism for the relatively high and consistent levels of self-recruitment documented for Kimbe Island.

Results obtained from our demographic model showed that the three subpopulations with the lowest mortality rates (A, B, and E) are capable of persisting without immigration. Two of these subpopulations (A and E) also have the largest anemone abundance and correspondingly have the highest number of clownfish and the highest self-recruitment rates. Yet, the female reproductive success for self-recruitment was quite different between subpopulations A and E despite their numbers differing by only 2%. The biannual production of self-recruiting juveniles per females from subpopulation A was more than twice that documented for E (approx. 0.38 versus approx. 0.16), perhaps due to local water movement, ability of larvae to sense the shallower anemones here, or a combination of these. Two different demographic strategies can ensure the persistence of subpopulations in the absence of immigration. The first strategy relies upon high self-recruitment to compensate for high mortality (subpopulation A). The second is the inverse; female mortality rates may be low so self-recruitment rates can be low as well (subpopulations B and E). In contrast to A, B, and E, subpopulations C, D, F, and G disappeared quickly without immigration. Subpopulation F consisted of only seven inhabited anemones in total. For the three other subpopulations (C, D, and G) persistence was not assured once isolated due to higher mortality rates and lower self-recruitment rates in comparison to subpopulations A, B, and E. The persistence of the island

409 population as a whole was driven by the low mortality rates of females and the high self-  
410 recruitment rates characteristic of the largest subpopulations at Kimbe Island (i.e. A, B, and E).

411 In conclusion, longitudinal genetic monitoring of each individual within a population is a powerful  
412 approach for understanding the local drivers of population persistence. It can provide accurate  
413 measurements of demographic rates that often cannot be reliably estimated by traditional  
414 approaches. Our use of longitudinal genetic monitoring is why we are confident it is possible for the  
415 Kimbe Island population to persist without immigration; i.e. that our primary result is robust. While  
416 we cannot be absolutely certain as to whether the Kimbe Island clownfish population will persist  
417 without immigration, all of the evidence presented suggests it is possible. We were also able to  
418 reliably determine the demographic parameters driving local-scale spatial variation in persistence  
419 likelihood, which, with further work, could be used to examine how and whether local adaptation  
420 develops and manifests. The approach outlined here will also be of help to managers in developing  
421 appropriately targeted and scaled actions that maximize persistence likelihood in this era of climate  
422 change and increasing disturbance frequencies. For example, our results indicate that protecting  
423 Kimbe Island may be sufficient to preserve the local *A. percula* population over time even without  
424 immigration from surrounding populations.

425 **Ethics statement.** The research accomplished in this project was conducted under James Cook  
426 University ethics approval number A1264 and followed all guidelines for the country in which it  
427 took place.

428 **Authors' contributions.** SP, GPJ, and SRT conceived the study; OCS, PSA, GRA, GPJ, MLB,  
429 SRT, and SP collected field data; OCS and PSA carried out the molecular lab work and parentage  
430 analysis, OCS, JAM, and SP drafted the manuscript; GRA, SRT, GPJ, and MLB helped to draft the  
431 manuscript; OCS carried out the statistical analyses; OCS, CMB and MJ developed and analyzed  
432 the demographic model. All authors gave final approval for publication.

**Acknowledgements.** We thank all those that assisted in the field. N. Tolou and S. Baksay assisted with genetic analysis. Mahonia Na Dari R and CC and the crew of M.V. FeBrina provided essential logistic support. We are grateful to the traditional owners of the reefs near Kimbe Island, PNG, for allowing us access to their reefs.

**Funding statement.** This research was supported by LABEX Corail, ERC, CRISP, the GEF CRTR Connectivity Working Group, NSF, the ARC CoE Coral Reef Studies, TNC, Total Foundation, JCU, KAUST, and WHOI.

## References

1. Levin SA. 1992 The problem of pattern and scale in ecology. *Ecology* 73, 1943–1967. (doi:10.2307/1941447)
2. Hastings A, Botsford LW. 2006 Persistence of spatial populations depends on returning home. *Proc. Natl Acad. Sci. USA* 103, 6067–6072. (doi:10.1073/pnas.0506651103)
3. Austin A, Ovaskainen O, Hanski I. 2011 Size and genetic composition of the colonizing propagules in a butterfly metapopulation. *Oikos* 120, 1357–1365. (doi:10.1111/j.1600-0706.2010.18992.x)
4. Hanski I. 1998 Metapopulation dynamics. *Nature* 396, 41–49. (doi:10.1038/23876)
5. Cowen RK, Sponaugle S. 2009 Larval dispersal and marine population connectivity. *Annu. Rev. Mar. Sci.* 1, 443–466. (doi:10.1146/annurev.marine.010908.163757)
6. White JW, Botsford LW, Hastings A, Largier JL. 2010 Population persistence in marine reserve networks: incorporating spatial heterogeneities in larval dispersal. *Mar. Ecol. Prog. Ser.* 398, 49–67. (doi:10.3354/meps08327)

- 454 7. Murdoch WW. 1994 Population regulation in theory and practice. *Ecology* 75, 271–287.  
455 (doi:10.2307/ 1939533)
- 456 8. Cappuccino N, Price PW. 1995 Population dynamics: new approaches and synthesis.  
457 Amsterdam, The Netherlands: Elsevier Science.
- 458 9. Gerber LR, Botsford LW, Hastings A, Possingham HP, Gaines SD, Palumbi SR, Andelman S.  
459 2003 Population models for marine reserve design: a retrospective and prospective synthesis. *Ecol.*  
460 *Appl.* 13, S47–S64. (doi:10.1890/1051- 0761(2003)013[0047:PMFMRD]2.0.CO;2)
- 461 10. Almany GR, Berumen ML, Thorrold SR, Planes S, Jones GP. 2007 Local replenishment of  
462 coral reef fish populations in a marine reserve. *Science* 316, 742–744.  
463 (doi:10.1126/science.1140597)
- 464 11. Hanski I. 1999 Metapopulation ecology. Oxford, UK: Oxford University Press.
- 465 12. Pollock KH. 1991 Modeling capture, recapture, and removal statistics for estimation of  
466 demographic parameters for fish and wildlife populations - past, present and future. *J. Amer. Stat.*  
467 *Asso.* 86, 225–238.
- 468 13. Sandercock BK. 2006 Estimation of demographic parameters from live-encounter data: a  
469 summary review. *J. Wildl. Manag.* 70, 1504–1520. (doi:10. 2193/0022-  
470 541X(2006)70[1504:EODPFL]2.0.CO;2)
- 471 14. Thompson W. 2013 Sampling rare or elusive species: concepts, designs, and techniques for  
472 estimating population parameters. Washington, USA: Island Press.
- 473 15. Burgess SC et al. 2014 Beyond connectivity: how empirical methods can quantify population  
474 persistence to improve marine protected-area design. *Ecol. Appl.* 24, 257–270. (doi:10.1890/13-  
475 0710.1)

- 476 16. Cain ML, Milligan BG, Strand AE. 2000 Long distance seed dispersal in plant populations.  
477 Amer. J. Bot. 87, 1217–1227. (doi:10.2307/ 2656714)
- 478 17. Schaub M, Reichlin TS, Abadi F, Kary M, Jenni L, Arlettaz R. 2012 The demographic drivers  
479 of local population dynamics in two rare migratory birds. *Oecology* 168, 97–108.  
480 (doi:10.1007/s00442-011-2070-5)
- 481 18. Pulliam HR. 1988 Sources, sinks, and population regulation. *Amer. Nat.* 132, 652–661.  
482 (doi:10.1086/ 284880)
- 483 19. Cowen RK, Lwiza KMM, Sponaugle S, Paris CB, Olson DB. 2000 Connectivity of marine  
484 populations: open or closed? *Science* 287, 857–859. (doi:10. 1126/science.287.5454.857)
- 485 20. Cowen RK, Paris CB, Srinivasan A. 2006 Scaling of connectivity in marine populations.  
486 *Science* 311, 522–527. (doi:10.1126/science.1122039)
- 487 21. Swearer SE, Caselle JE, Lea DW, Warner RR. 1999 Larval retention and recruitment in an  
488 island population of a coral-reef fish. *Nature* 402, 799– 802. (doi:10.1038/45533)
- 489 22. Carson HS, Cook GS, Lopez-Duarte PC, Levin LA. 2011 Evaluating the importance of  
490 demographic connectivity in a marine metapopulation. *Ecology* 92, 1972–1984. (doi:10.1890/11-  
491 0488.1)
- 492 23. Gilg MR, Hilbish TJ. 2003 The geography of marine larval dispersal: coupling genetics with  
493 fine-scale physical oceanography. *Ecology* 84, 2989–2998. (doi:10.1890/02-0498)
- 494 24. Planes S, Jones GP, Thorrold SR. 2009 Larval dispersal connects fish populations in a network  
495 of marine protected areas. *Proc. Natl Acad. Sci. USA* 106, 5693–5697.  
496 (doi:10.1073/pnas.0808007106)

- 497 25. Berumen ML, Almany GR, Planes S, Jones GP, Saenz-Agudelo P, Thorrold SR. 2012  
498 Persistence of self-recruitment and patterns of larval connectivity in a marine protected area  
499 network. *Ecol. Evol.* 2, 444–452. (doi:10.1002/ece3.208)
- 500 26. Buston PM. 2003 Mortality is associated with social rank in the clown anemonefish  
501 (*Amphiprion percula*). *Mar. Biol.* 143, 811–815. (doi:10.1007/ s00227-003-1106-8)
- 502 27. Allen GR. 1975 The anemonefishes: their classification and biology. California, USA: T.F.H.  
503 Publications.
- 504 28. Fautin DG, Allen GR. 1997 Anemone fishes and their host sea anemones: a guide for aquarists  
505 and divers. Perth, Australia: Western Australian Museum.
- 506 29. Buston P. 2004 Does the presence of non-breeders enhance the fitness of breeders? An  
507 experimental analysis in the clown anemonefish *Amphiprion percula*. *Behav. Ecol. Sociobiol.* 57,  
508 23–31. (doi:10. 1007/s00265-004-0833-2)
- 509 30. Fricke HW. 1979 Mating system, resource defense and sex change in the anemonefish  
510 *Amphiprion akallopisos*. *Z. Tierphysiol. Tierer.* 50, 313–326.
- 511 31. Hattori A. 1994 Intergroup movement and mate acquisition tactics of the protandrous  
512 anemonefish, *Amphiprion clarkii*, on coral-reef, Okinawa. *Jpn. J. Ichthyol.* 41, 159–165.
- 513 32. Buston P. 2003 Forcible eviction and prevention of recruitment in the clown anemonefish.  
514 *Behav. Ecol.* 14, 576–582. (doi:10.1093/beheco/arg036)
- 515 33. Mitchell JS. 2003 Social correlates of reproductive success in false clown anemonefish:  
516 subordinate group members do not pay-to-stay. *Evol. Ecol. Res.* 5, 89–104.


34. Berumen ML, Walsh HJ, Raventos N, Planes S, Jones GP, Starczak V, Thorrold SR. 2010 Otolith geochemistry does not reflect dispersal history of clownfish larvae. *Coral Reefs* 29, 883–891. (doi:10.1007/s00338-010-0652-z)
35. Peakall R, Smouse PE. 2012 GenAlEx 6.5: genetic analysis in Excel. Population genetic software for teaching and research—an update. *Bioinformatics* (Oxford, England) 28, 2537–2539. (doi:10.1093/bioinformatics/bts460)
36. Saenz-Agudelo P, Jones GP, Thorrold SR, Planes S. 2009 Estimating connectivity in marine populations: an empirical evaluation of assignment tests and parentage analysis under different gene flow scenarios. *Mol. Ecol.* 18, 1765–1776. (doi:10.1111/j.1365-294X.2009.04109.x)
37. Gerber S, Chabrier P, Kremer A. 2003 FAMOZ: a software for parentage analysis using dominant, co-dominant and uni-parentally inherited markers. *Mol. Ecol. Notes* 3, 479–481. (doi:10.1046/j.1471-8286.2003.00439.x)
38. R-Development-Core-Team. 2007 R: a language and environment for statistical computing., ed computing RfCs Vienna, Austria.
39. Hofbauer J, Sigmund K. 1998 Evolutionary games and population dynamics. Cambridge, UK: Cambridge University Press.
40. Soetaert K, Herman PMJ. 2008 A practical guide to ecological modelling: using R as a simulation platform. Berlin, Germany: Springer.
41. Nelder JA, Mead R. 1965 A simple method for function minimization. *Comput. J.* 7, 308–313. (doi:10.1093/comjnl/7.4.308)
42. Nichols JD, Pollock KH. 1990 Estimation of recruitment from immigration versus in situ reproduction using Pollock's robust design. *Ecology* 71, 21–26. (doi:10.2307/1940243)

- 539 43. Kendall WL, Nichols JD, Hines JE. 1997 Estimating temporary emigration using capture-  
540 recapture data with Pollock's robust design. *Ecology* 78, 563–578.
- 541 44. Pine WE, Pollock KH, Hightower JE, Kwak TJ, Rice JA. 2003 A review of tagging methods for  
542 estimating fish population size and components of mortality. *Fisheries* 28, 10–23.  
543 (doi:10.1577/1548- 8446(2003)28[10:AROTMF]2.0.CO;2)
- 544 45. Kininmonth S et al. 2011 Dispersal connectivity and  
545 reserve selection for marine conservation. *Ecol.*  
546 *Model.* 222, 1272–1282. (doi:10.1016/j.ecolmodel. 2011.01.012)
- 547 46. Hughes TP, Baird AH, Dinsdale EA, Moltschaniwskyj NA, Pratchett MS, Tanner JE, Willis  
548 BL. 1999 Patterns of recruitment and abundance of corals along the Great Barrier Reef. *Nature* 397,  
549 59–63. (doi:10.1038/16237)
- 550 47. Hogan JD, Thiessen RJ, Heath DD. 2010 Variability in connectivity indicated by chaotic  
551 genetic patchiness within and among populations of a marine fish. *Mar. Ecol. Prog. Ser.* 417,  
552 U263–U289. (doi:10.3354/ meps08793)
- 553 48. Leis JM, Carson-Ewart BM, Hay AC, Cato DH. 2003 Coral-reef sounds enable nocturnal  
554 navigation by some reef-fish larvae in some places and at some times. *J. Fish Biol.* 63, 724–737.  
555 (doi:10.1046/j. 1095-8649.2003.00182.x)
- 556 49. Simpson SD, Meekan M, Montgomery J, McCauley R, Jeffs A. 2005 Homeward sound.  
557 *Science* 308, 221. (doi:10.1126/science.1107406)
- 558 50. Montgomery JC, Jeffs A, Simpson SD, Meekan M, Tindle C. 2006 Sound as an orientation cue  
559 for the pelagic larvae of reef fishes and decapod crustaceans. In *Advances in marine biology*, vol.  
560 51 (eds AJ Southward, DW Sims), pp. 143–196. San Diego, USA: Academic Press.




- 561 51. Atema J, Kingsford MJ, Gerlach G. 2002 Larval reef fish could use odor for detection, retention  
562 and orientation to reefs. *Mar. Ecol. Prog. Ser.* 241, 151–160. (doi:10.3354/meps241151)
- 563 52. Doving KB, Stabell OB, Ostlund-Nilsson S, Fisher R. 2006 Site fidelity and homing in tropical  
564 coral reef cardinalfish: are they using olfactory cues? *Chem. Senses* 31, 265–272.  
565 (doi:10.1093/chemse/bjj028)
- 566 53. Gerlach G, Atema J, Kingsford MJ, Black KP, Miller- Sims V. 2007 Smelling home can  
567 prevent dispersal of reef fish larvae. *Proc. Natl Acad. Sci. USA* 104, 858–863.  
568 (doi:10.1073/pnas.0606777104)
- 569 54. Cowen RK, Gawarkiewicz G, Pineda J, Thorrold SR, Werner FE. 2007 Population connectivity  
570 in marine systems an overview. *Oceanography* 20, 14–21. (doi:10.5670/oceanog.2007.26)
- 571 55. Thorrold SR, Zacherl DC, Levin LA. 2007 Population connectivity and larval dispersal using  
572 geochemical signatures in calcified structures. *Oceanography* 20, 80–89.  
573 (doi:10.5670/oceanog.2007.31)
- 574 56. Dixon DL, Jones GP, Munday PL, Planes S, Pratchett MS, Srinivasan M, Syms C, Thorrold  
575 SR. 2008 Coral reef fish smell leaves to find island homes. *Proc. R. Soc. B* 275, 2831–2839.  
576 (doi:10.1098/rspb.2008.0876)

577 **Figure captions**



578

579 **Figure 1.** Self-recruitment (SR, blue arrows), local connectivity (LC, grey arrows), and  
580 immigration (I, black arrows) calculated on the natural *Amphiprion percula* populations from  
581 parentage analysis from 2005 to 2013 in Kimbe Island, Papua New Guinea. Subpopulations are  
582 indicated with letters A-G. The numbers correspond to new-recruits and the arrows show the  
583 sources of these inputs (i.e. self-recruitment, local connectivity, or immigration).



584

585 **Figure 2.** Demographic model projections of numbers of juvenile *Amphiprion percula* within each of the subpopulations (A - G) and the  
 586 population of Kimbe Island (subpopulations A-G combined) for three different recruitment and immigration scenarios. The model was initialized  
 587 with data of total juvenile captures during 2007 (set as time[year] = 0). The model temporal resolution is one month and the results are plotted here  
 588 every six months for 50 years. Self-recruitment and local connectivity refer to reproductive success to self-recruitment and reproductive success to  
 589 local connectivity, respectively, in the demographic model.

590 **Table**591 **Table 1.** Demographic parameters from empirical values of natural population of *A. percula*.

592  $d_{J,M,F}^i$  refers to average biannual mortality rate per capita of juvenile, and adult males and females  
 593 for each subpopulation between two periods 2009-2011 and 2011-2013 (see Table S1 for mortality  
 594 rates for each of the two 2-year periods).  $n^i$  is the maximal average number of juvenile per  
 595 anemone in the subpopulation  $i$  between 2007 and 2013 and  $K^i$  is the total number of anemones in  
 596 the subpopulation  $i$  (equal to the maximal number of male or female per subpopulation).

| Parameters | A     | B     | C     | D     | E     | F     | G     |
|------------|-------|-------|-------|-------|-------|-------|-------|
| $d_J^i$    | 0.296 | 0.270 | 0.269 | 0.477 | 0.179 | 0.500 | 0.487 |
| $d_M^i$    | 0.373 | 0.095 | 0.393 | 0.400 | 0.198 | 0.286 | 0.440 |
| $d_F^i$    | 0.397 | 0.190 | 0.382 | 0.350 | 0.243 | 0.357 | 0.552 |
| $n^i$      | 2.19  | 2.00  | 1.58  | 1.85  | 2.03  | 1.14  | 1.70  |
| $K^i$      | 57    | 21    | 19    | 20    | 56    | 7     | 30    |

597